




Rheumatoid Arthritis



Dr. Majid Ghayour Mobarhan

Nutritionist from the UK

Rheumatoid arthritis signs and symptoms may vary in severity and may even come and go. Periods of increased disease activity, called flares, alternate with periods of relative remission - when the swelling and pain fade or disappear. Over time, rheumatoid arthritis can cause joints to deform and shift out of place.

When to see a doctor:

Make an appointment with your doctor if you have persistent discomfort and swelling in your joints.

➤ Causes ◀

Rheumatoid arthritis occurs when your immune system attacks the synovium - the lining of the membranes that surround your joints.

The resulting inflammation thickens the synovium, which can eventually destroy the cartilage and bone within the joint.

The tendons and ligaments that hold the joint together weaken and stretch. Gradually, the joint loses its shape and alignment.

Doctors don't know what starts this process, although a genetic component appears likely.

While your genes don't actually cause rheumatoid arthritis, they can make you more susceptible to environmental factors - such as infection with certain viruses and bacteria - that may trigger the disease.

➤ Risk factors ◀

Factors that may increase your risk of rheumatoid arthritis include:

- ➤ Your sex: Women are more likely than men to develop rheumatoid arthritis.
 - Age: Rheumatoid arthritis can occur at any age, but it most commonly begins in middle age.



- Family history: If a member of your family has rheumatoid arthritis, you may have an increased risk of the disease.
- risk of developing rheumatoid arthritis, particularly if you have a genetic predisposition for developing the disease. Smoking also appears to be associated with greater disease severity.
- ▶ Environmental exposures: Although poorly understood, some exposures such as asbestos or silica may increase the risk of developing rheumatoid arthritis. Emergency workers exposed to dust from the collapse of the World Trade Center are at higher risk of autoimmune diseases such as rheumatoid arthritis.
- Obesity: People especially women age 55 and younger who are overweight or obese appear to be at a somewhat higher risk of developing rheumatoid arthritis.



Rheumatoid arthritis is a chronic inflammatory disorder that can affect more than just your joints. In some people, the condition can damage a wide variety of body systems, including the skin, eyes, lungs, heart and blood vessels.

As an autoimmune disorder, rheumatoid arthritis occurs when your immune system mistakenly attacks your own body's tissues.

Unlike the wear-and-tear damage of osteoarthritis, rheumatoid arthritis affects the lining of your joints, causing a painful swelling that can eventually result in bone erosion and joint deformity.

The inflammation associated with rheumatoid arthritis is what can damage other parts of the body as well. While new types of medications have improved treatment options dramatically, severe rheumatoid arthritis can still cause physical disabilities.



Signs and symptoms of rheumatoid arthritis may include:

- ▶ Tender, warm, swollen joints
- ➤ Joint stiffness that is usually worse in the mornings and after inactivity
 - Fatigue, fever and loss of appetite



Early rheumatoid arthritis tends to affect your smaller joints first - particularly the joints that

attach your fingers to your hands and your toes to your feet.

