

# The Effect of Maternal Selenium Supplementation on Pregnancy Outcome and the Level of Oxidative Stress in Neonates

H Boskabadi<sup>1</sup>, F Rezagholizade Omran<sup>1</sup>, F Tara<sup>2</sup>, MP Rayman<sup>3</sup>, M Ghayour-Mobarhan<sup>4,5\*</sup>, A Sahebkar<sup>4</sup>, S Tavallaie<sup>4</sup>, MT Shakeri<sup>6</sup>, DH Alamdari<sup>5</sup>, M Kiani<sup>7</sup>, BS Razavi<sup>4</sup>, M Oladi<sup>4</sup>, G Ferns<sup>8</sup>

<sup>1</sup>Neonatal Research Center, <sup>2</sup>Department of Obstetrics and Gynecology, OM-Albanin Hospital, Mashhad University of Medical Sciences, Mashhad, Iran, <sup>3</sup>Division of Nutrition Science, Faculty of Health, School of Biomedical and Molecular Sciences, University of Surrey, Guildford GU2 7XH, UK, <sup>4</sup>Cardiovascular Research Center, Avicenna Research Institute, <sup>5</sup>Department of Biochemistry and Nutrition, <sup>6</sup>Department of Statistics, <sup>7</sup>Department of Obstetrics and Gynecology, Qaem Hospital, Mashhad University of Medical Sciences, Mashhad, Iran, <sup>8</sup>Institute for Science and Technology in Medicine, University of Keele, Guy Hilton Research Centre, UK

## Abstract

**Background:** Pregnancy is regarded as a condition which is usually accompanied by oxidative stress. This study was undertaken to investigate the effect of maternal selenium supplementation during gestation on the level of oxidative stress in neonates and the pregnancy outcome.

**Methods:** In this double-blind trial, 179 primigravid pregnant women in the first trimester of pregnancy were randomly assigned to receive 100 µg of selenium (Se group) or a placebo (control group) per day until delivery. The level of oxidative stress and serum selenium concentration was determined in the maternal and umbilical cord sera of the subjects. Oxidative stress was measured by means of a novel assay of prooxidant-antioxidant balance (PAB). The incidence of any pregnancy complications and outcomes was also evaluated in all neonates, being fully examined and followed up until 45 days.

**Results:** Although maternal selenium concentration was significantly higher in the Se group ( $p < 0.001$ ), there was no statistically significant differences in the umbilical cord selenium content between the two groups. Selenium supplementation was not associated with any significant decrease in PAB values in the Se group. The incidence of neonatal complications and outcomes did not differ significantly between the groups.

**Conclusion:** Maternal selenium supplementation during pregnancy was safe but was not associated with a significant change in the extent of oxidative stress in neonates.

**Keywords:** Selenium; Prooxidant-Antioxidant balance; Neonate; Complication; Pregnancy

## Introduction

Oxidative stress is defined as an imbalance between the generation of reactive oxygen species (ROS) and

the ability of biological antioxidant systems to neutralize these species.<sup>1</sup> Pregnancy is regarded as a condition which is usually accompanied by oxidative stress.<sup>2,3</sup> This is due to the physiological changes that occur during pregnancy including the increase in the basal oxygen intake and consumption as well as elevated metabolic demand, finally leading to the overproduction of reactive oxygen species and decreased activity of antioxidant enzymes.<sup>4,5</sup> On the other

\*Correspondence: Majid Ghayour-Mobarhan, MD, Department of Biochemistry and Nutrition, Faculty of Medicine, Mashhad University of Medical Science, Mashhad, Iran. Tel: +98-511-8828573, Fax: +98-511-8828574, e-mail: ghayourm@mums.ac.ir  
Received: July 10, 2009 Accepted: November 10, 2009

hand, labor is also a very stressful condition which is itself a cause of significant oxidative stress both in the mother and infant.<sup>6</sup> Moreover, other mechanisms such as exposure to a high oxygen concentration at birth, infection, inflammation and a deficient antioxidant system make the newborn more susceptible to oxidative stress.<sup>7,8</sup> Reactive oxygen species are considered to have a role in the pathophysiology of various neonatal diseases such as bronchopulmonary dysplasia, retinopathy of prematurity, persistent ductus arteriosus, narcotizing enterocolitis, intracranial hemorrhage and hypoxic ischemic encephalopathy.<sup>9-12</sup>

According to the literature, the risk of oxidative stress in neonates depends largely on the maternal antioxidant status which is potentially important for the protection of maternal-fetal unit against free radicals.<sup>3,13,14</sup> As previously stated, pregnancy is considered as a condition that favors oxidative stress and during pregnancy, lower concentrations of antioxidant enzymes have been reported.<sup>2-5</sup> Regarding the significant correlation existing between oxidative status of the mother and neonate,<sup>15</sup> one of the effective strategies against the development of oxidative stress in the newborn is the enhancement of maternal antioxidant system via supplementation with antioxidant compounds.

Selenium is an essential trace micronutrient element which functions in the human body as a key component of some important selenoproteins including the antioxidant enzymes glutathione peroxidases and thioredoxin reductases.<sup>16</sup> Compared with pre-pregnancy levels, lower concentrations of selenium in the whole blood and plasma as well as lower glutathione peroxidase activities have been reported in pregnant women.<sup>17,18</sup> The same trend exists in non-pregnant controls.<sup>19</sup> This decrease in selenium status which is progressive as gestation proceeds may be partly attributed to hemodilution from the blood volume increase associated with pregnancy.<sup>17,19</sup>

In the current study, we sought to investigate the impact of selenium supplementation in pregnant women on the extent of oxidative stress in their newborn. For the evaluation of oxidative stress, we applied a recently developed and modified prooxidant-antioxidant balance (PAB) assay which is a simple, rapid and inexpensive test.<sup>20,21</sup> Furthermore, we looked into the effect of selenium supplementation on the incidence any complications to assess the safety of maternal selenium supplementation during pregnancy for the newborn.

## Materials and Methods

Two hundred and eighteen pregnant women with an age range of 16-35 years were assessed for eligibility to participate in this trial. The subjects were selected from women referring to the Obstetrics and Gynecology Department of OM-Albanin Hospital (Mashhad, Iran) between June 2006 and August 2008. The inclusion criteria for selection were gestational age up to 12 weeks with no indications for terminating the pregnancy. Exclusion criteria included the use of any drugs, except routine supplements of folic acid and ferrous sulfate, and a prior history or clinical features of any medical conditions, including thyroid disorders, diabetes, hypertension and infections. Thirty nine individuals were excluded from the study with 179 subjects entering the trial. Of these individuals, 13 were excluded because of intolerance to ( $n=4$ ) or the unpleasant aroma associated with the tablets ( $n=9$ ). In a double-blind manner, the 166 eligible subjects were randomized to receive 100  $\mu\text{g}$  of selenium, as selenium yeast, daily from the first trimester of their pregnancy until delivery, a period of approximately six months (Se group,  $n=83$ ), or daily placebo yeast tablets for the same period (control group,  $n=83$ ). One hundred and twenty-five subjects completed the study ( $n=61$  and 64 for Se and Control groups, respectively). Drop-outs ( $n=41$ ) mainly comprised those who were ultimately reluctant or unable to comply with the regular consumption of medication (selenium or placebo), or who suffered a miscarriage.

The neonates included in the study were all those who were delivered from participant mothers. For each newborn, a sequence of examinations was performed by a neonatologist in the three intervals of 3-5, 10-15 and 30-45 days of birth. Each subject gave written consent to participate in the study. The study protocol was approved by the Ethics Committee for Clinical Research of Mashhad University of Medical Sciences (MUMS).

For all the newborns, the required parameters including gender, weight, length, gestational age, head circumference and Apgar scores in the first and fifth minutes of birth were determined. Head circumference was measured at the level of occipital protuberance and frontal. The body weight of each subject was measured with a standard scale to an accuracy of 0.1 gram. The Apgar score was evaluated in the first and fifth minutes of birth in the labor room. The gestational age was measured in terms of weeks from the last menstrual period and accordingly an immature newborn was defined as a newborn with

gestational age < 37 weeks and a mature newborn as a newborn with gestational age ≥ 37 weeks.

The newborn complications which were evaluated included newborn mortality, immaturity, pathologic icterous, birth asphyxia, intraventricular hemorrhage (IVH), Apgar score at the fifth minutes < 7, and respiratory distress syndrome. The newborn anomalies evaluated included hydrocephaly, torticollis, fetal dent, hypospadiasis, mandibular hypoplasia, imperforated anus, valgus metatars, and developmental dysplasia of the hip.

Maternal and umbilical cord blood samples were collected after birth and centrifuged at 2500 rpm for 15 min at room temperature to obtain the serum. Hemolyzed samples were excluded from analysis. Serum was stored at -20°C prior to analysis.

Serum selenium was determined by electrothermal atomic absorption spectrometry with Zeeman background correction using a palladium chloride chemical modifier.<sup>22,23</sup>

To assess the oxidative stress level in the newborns, a modified PAB assay was applied based on a previously described method.<sup>20,21</sup>

All statistical analyses were performed using SPSS software (version 11.5, Chicago, IL, USA). The values were expressed as mean ± SD (for normally distributed

data) or median and interquartile range (for non-normally distributed data). The group comparisons were carried out using *t* test (for normally distributed data) or Mann-Whitney U test (for non-normally distributed data). Categorical data were compared by Fisher's Exact test. A two-tailed *p*-value of < 0.05 was considered as statistically significant.

## Results

### *Effect of Selenium Supplementation on the Incidence of Complications in Newborns*

Newborn parameters including gender, gestational age, birth weight, birth length, head circumference and Apgar scores at the first and the fifth minutes of birth did not differ significantly between the Se and control groups (*p* > 0.05, Table 1). There was also no significant difference in newborn complications (mortality, immaturity, pathologic icterous, asphyxia, intraventricular hemorrhage (IVH), Apgar score at the fifth minutes < 7 and respiratory distress syndrome) and anomalies (hydrocephaly, torticollis, fetal dent, hypospadiasis, mandibular hypoplasia, imperforated anus, valgus metatars and developmental dysplasia of the hip) between the newborns in the Se and control

**Table 1:** Characteristics of neonates in the Se and control groups.

Parameter	Se group	Control group	P value
Male (%)	55.8	58.2	0.85
Gestational age (weeks)	39.0 (39.0-40.0)	39.0 (38.0-40.0)	0.97
Birth weight (g)	3085.3 ± 622.2	3069.0 ± 551.1	0.88
Birth length (cm)	50.0 (49.0-51.0)	50.0 (48.0-51.0)	0.76
Birth head circumflax (cm)	34.4 ± 1.4	34.5 ± 2.9	0.53
Apgar at 1 min (n)	9.0 (8.0-9.0)	9.0 (8.0-9.0)	0.63
Apgar at 5 min (n)	9.0 (9.0-9.0)	9.0 (9.0-9.0)	0.66

Values are expressed as percentile, mean ± SD or median and interquartile range.

**Table 2:** Comparison of the occurrences of neonatal complications and anomalies between Se and control groups.

Parameter	Se group (%)	Control group (%)	P value	
Mortality	Congenital anomaly	0	1.6	1.00
	Infection	1.9	0	0.45
	Immaturity	1.9	1.6	1.00
	Immaturity	7.7	9.2	1.00
	Pathologic icter	1.7	1.6	1.00
	Birth asphyxia	7.7	7.9	1.00
Morbidity	Respiratory distress syndrome	8.3	9.5	1.00
	IVH	0	3.1	0.50
	Apgar < 7	3.4	4.7	1.00
	Congenital anomaly	3.8	6.0	0.69

IVH: Intraventricular hemorrhage.

groups ( $p>0.05$ , Table 2).

In spite of the significantly higher maternal serum selenium concentration in the Se group compared with that in the control group ( $p<0.001$ , Table 3), there was no significant difference in the umbilical cord serum selenium levels between the two groups ( $p=0.294$ , Table 3). Supplementation with selenium was not associated with a significant decrease in the umbilical cord PAB values in the Se group compared with that in the control group ( $p=0.190$ , Table 3).

### Discussion

In the present study, we found no significant effect of selenium supplementation (100 µg/day) on neonatal parameters and outcomes during the postpartum 45-day follow-up period. In a previous trial among 913 HIV-infected pregnant women, it was reported that selenium supplementation (200 µg/day) was not associated with a significant effect on pregnancy outcomes, neonatal parameters and mortality, but it reduced the risk of child mortality after 6 weeks. These findings were based on a longer follow-up (until 6 months after delivery) and larger supplementation dose (200 µg/day).<sup>22</sup> In another trial, postnatal selenium supplementation in very low birth weight infants was not found to improve neonatal outcomes.<sup>23</sup>

In spite of the significantly higher maternal serum selenium concentrations in the supplemented vs. non-supplemented group ( $p<0.001$ ), there was no significance difference in cord selenium levels between the groups ( $p>0.05$ ). To the best of our knowledge, to date there has been no report about the level of selenium in Iranian neonates. The mean selenium concentration which was measured in the cord blood of the subjects was higher than that of those in countries like Germany, Finland, Slovenia and Poland,<sup>24-27</sup> while being lower than that of those in Japan, Turkey and California in the USA.<sup>28-30</sup> These differences in selenium status are probably due to the regional and geographical variability in the selenium content of

soil and plant foods because selenium enters food chain primarily through plants (as reviewed by Rayman).<sup>31</sup> The cord blood selenium in our study was approximately 63% of the maternal selenium. In contrast to some studies,<sup>18,29,27</sup> our findings is consistent with the majority of previous reports on the concentration of selenium in maternal and cord blood.<sup>26,32-35</sup> Lower levels of cord selenium compared to maternal selenium concentrations might be attributed to the high metabolic demands by the fetal tissues or the action of metallothionein-1. Metallothionein-1 is a heavy naturally occurring metal-binding protein in human placenta that sequesters some metals including selenium and can influence the distribution of selenium between serum and tissues. However, other mechanisms such as limited transport of selenium across the placenta may also be responsible for the lower cord selenium concentration.<sup>35</sup>

Regarding the good correlation existing between the maternal and cord selenium concentration,<sup>15</sup> maternal selenium deficiency is considered to be a contributory factor in the causation of oxidative stress in the neonate which is characterized by increased lipid peroxidation and impaired development of the immune system in the neonate.<sup>36,37</sup> However, the findings of our study indicate that selenium supplementation during gestation does not lead to a significant change in the extent of oxidative stress in the neonate. Our results also showed that there was no significant difference in the incidence of neonatal anomalies and outcomes between the groups. This major finding confirms that selenium supplementation during pregnancy is safe and is not associated with any side effect for the neonate. This safety for the neonate is especially important for the mothers who may wish to benefit from selenium supplementation during pregnancy and protect themselves against the heightened state of oxidative stress and decreased selenium status that occur during pregnancy.<sup>2-5,17-19</sup>

The method applied here for the estimation of oxidative stress was a modified PAB assay which has been recently developed by our group.<sup>21</sup> Briefly, in this method total prooxidant and antioxidant capacities are measured in a single assay and a redox index

**Table 3:** Comparison of maternal and umbilical cord selenium concentrations and PAB values between Se and control groups

Parameter	Se group	Control group	P value
Maternal selenium (µg/L)	168.6±36.4	119.4±33.4	< 0.001
Cord selenium (µg/L)	106.3±18.2	101.9±15.9	0.29
PAB (AU)	37.2 (26.1-121.0)	30.8 (24.0-45.5)	0.19

Values are expressed as mean±SD or median and interquartile range. PAB: prooxidant-antioxidant balance. AU: arbitrary unit.

is obtained. This method has been calibrated against a series of well-known oxidants and antioxidants and its response has been found to be linear.<sup>20</sup> Finally, the simple, rapid and inexpensive procedure of this method poses the probability of its application as a routine clinical laboratory test for the assessment of oxidative stress in neonates, who are extremely vulnerable to free radical damage, and also the choice of an effective prophylaxis for those who are at an increased risk of developing oxidative stress.

The major limitation of the current study was the duration of follow-up period. With a longer follow-up, we would be able to obtain a better assessment of the impact of selenium supplementation on neonatal abnormalities and outcomes that may occur later than 45 days of life. Besides, determination of selenoprotein levels such as selenoprotein P could have been helpful for a better understanding of selenium status in both mothers and neonates.

In summary, the findings of our 45-day follow-up indicated that maternal selenium supplementation during pregnancy was safe for the newborns but was not

associated with a lower risk of oxidative stress. The results also suggest that the modified PAB assay might be useful (e.g. as a screening test) for the evaluation of oxidative status in the newborns. However, further and larger clinical research in various physiological and pathological states related to oxidative stress, and especially in pregnant women, who are selenium deficient, is required to confirm the feasibility and potency of this method as a clinical laboratory test.

### Acknowledgement

This work was supported by a grant from the Research Council of the Mashhad University of Medical Sciences (MUMS). The authors would also like to express their sincerest thanks to the staff of the Obstetrics and Gynecology Department of the OM-Albanin Hospital as well as Pharma Nord Vejle Company for their contribution to the study.

**Conflict of interest:** None declared.

### References

- Rice-Evans C, Burdon R. Free radical-lipid interactions and their pathological consequences. *Prog Lipid Res* 1993;**32**:71-110. [8415800] [doi:10.1016/0163-7827(93)90006-1]
- Chen X, Scholl TO. Oxidative stress: changes in pregnancy and with gestational diabetes mellitus. *Curr Diab Rep* 2005;**5**:282-8. [16033680] [doi:10.1007/s11892-005-0024-1]
- Arikan S, Konukoglu D, Arikan C, Akcay T, Davas I. Lipid peroxidation and antioxidant status in maternal and cord blood. *Gynecol Obstet Invest* 2001;**51**:145-9. [11306898] [doi:10.1159/000052913]
- Casanueva E, Viteri FE. Iron and oxidative stress in pregnancy. *J Nutr* 2003;**133**:1700S-1708S. [12730487]
- Walsh SW, Wang Y. Secretion of lipid peroxides by the human placenta. *Am J Obstet Gynecol* 1993;**169**:1462-6. [8267047]
- Rogers MS, Mongelli M, Tsang KH, Wang CC. Fetal and maternal levels of lipid peroxides in term pregnancies. *Acta Obstet Gynecol Scand* 1999;**78**:120-4. [10023874] [doi:10.1080/j.1600-0412.1999.780209.x]
- Gitto E, Pellegrino S, Gitto P, Barberi I, Reiter RJ. Oxidative stress of the newborn in the pre- and post-natal period and the clinical utility of melatonin. *J Pineal Res* 2009;**46**:128-39. [19054296] [doi:10.1111/j.1600-079X.2008.00649.x]
- Saugstad OD. Update on oxygen radical disease in neonatology. *Curr Opin Obstet Gynecol* 2001;**13**:147-53. [11315869] [doi:10.1097/00001703-200104000-00009]
- Buonocore G, Perrone S. Biomarkers of hypoxic brain injury in the neonate. *Clin Perinatol* 2004;**31**:107-16. [15183660] [doi:10.1016/j.clp.2004.03.008]
- Perrone S, Bracci R, Buonocore G. New biomarkers of fetal-neonatal hypoxic stress. *Acta Paediatr Suppl* 2002;**91**:135-8. [12477278] [doi:10.1080/080352502320764337]
- Drury JA, Nycyk JA, Baines M, Cooke RW. Does total antioxidant status relate to outcome in very pre-term infants? *Clin Sci (Lond)* 1998;**94**:197-201. [9536929]
- Saugstad OD. Oxidative stress in the newborn--a 30-year perspective. *Biol Neonate* 2005;**88**:228-36. [16210845] [doi:10.1159/000087586]
- Scholl TO, Chen X, Sims M, Stein TP. Vitamin E: maternal concentrations are associated with fetal growth. *Am J Clin Nutr* 2006;**84**:1442-8. [17158428]
- Buonocore G, Perrone S. Biomarkers of oxidative stress in the fetus and newborn. *Hematology* 2006;**2**:103-7.
- Argüelles S, Machado MJ, Ayala A, Machado A, Hervías B. Correlation between circulating biomarkers of oxidative stress of maternal and umbilical cord blood at birth. *Free Radic Res* 2006;**40**:565-70. [16753833] [doi:10.1080/10715760500519834]
- Letavayová L, Vlcková V, Brozmanová J. Selenium: from cancer prevention to DNA damage. *Toxicology* 2006;**227**:1-14. [16935405] [doi:10.1016/j.tox.2006.07.017]
- Zachara BA, Wardak C, Didkowskii W, Maciag A, Marchaluk E. Changes in blood selenium and glutathione concentrations and glutathione peroxidase activity in human pregnancy. *Gynecol Obstet Invest* 1993;**35**:12-7. [8449427] [doi:10.1159/000292655]
- Mihailović M, Cvetković M, Ljubić A, Kosanović M, Nedeljković S, Jovanović I, Pesut O. Selenium and malondialdehyde content and glutathione peroxidase activity in maternal and umbilical cord blood and amniotic fluid. *Biol Trace Elem Res* 2000;**73**:47-54. [10949968] [doi:10.1385/BTER:73:1:47]
- Ferrer E, Alegria A, Barbera R, Farre R, Lagarda MJ, Monleon J. Whole blood selenium content in



- pregnant women. *Sci Total Environ* 1999;**227**:139-43. [10231979] [doi:10.1016/S0048-9697(99)00008-X]
- 20 Alamdari DH, Paletas K, Pegiou T, Sarigianni M, Befani C, Koliakos G. Novel assay for the evaluation of the prooxidant-antioxidant balance, before and after antioxidant vitamin administration in type II diabetes patients. *Clin Biochem* 2007;**40**:248-54. [17196578] [doi:10.1016/j.clinbiochem.2006.10.017]
- 21 Alamdari DH, Ghayour-Mobarhan M, Tavallaie S, Parizadeh MR, Mohebbati M, Ghafoori F, Kazemi-Bajestani SM, Paletas K, Pegious T, Koliakos G. Prooxidant-antioxidant balance as a new risk factor in patients with angiographically defined coronary artery disease. *Clin Biochem* 2008;**41**:375-80. [18191639] [doi:10.1016/j.clinbiochem.2007.12.008]
- 22 Kupka R, Mugusi F, Aboud S, Msamanga GI, Finkelstein JL, Spiegelman D, Fawzi WW. Randomized, double-blind, placebo-controlled trial of selenium supplements among HIV-infected pregnant women in Tanzania: Effects on maternal and child outcomes. *Am J Clin Nutr* 2008;**87**:1802-8. [18541571]
- 23 Darlow BA, Winterbourn CC, Inder TE, Graham PJ, Harding JE, Weston PJ, Austin NC, Elder DE, Mo-gridge N, Buss IH, Sluis KB. The effect of selenium supplementation on outcome in very low birth weight infants: A randomized controlled trial. *J Pediatr* 2000;**136**:473-80. [10753245] [doi:10.1016/S0022-3476(00)90010-6]
- 24 Scramel P, Hasse S, Ovcar-Pavlu J. Selenium, cadmium, lead and mercury concentrations in human breast milk, in placenta, maternal blood, and the blood of the newborn. *Biol Trace Elem Res* 1988;**15**:111-24. [2484509] [doi:10.1007/BF02990130]
- 25 Dobrzynski W, Trafikowska U, Trafikowska A, Pilecki A, Szymanski W, Zachara BA. Decreased selenium concentration in maternal and cord blood in preterm compared with term delivery. *Analyst* 1998;**123**:93-7. [9581028] [doi:10.1039/a704884j]
- 26 Micetic-Turk D, Rossipal E, Krachler M, Li F. Maternal selenium status in slovenia and its impact on the selenium concentration of umbilical cord serum and colostrum. *Eur J Clin Nutr* 2000;**54**:522-4. [10878656] [doi:10.1038/sj.ejcn.1601050]
- 27 Korpela H, Loueniva R, Yrjänheikki E, Kauppila A. Selenium concentration in maternal and umbilical cord blood, placenta and amniotic membranes. *Int J Vitam Nutr Res* 1984;**54**:257-61. [6500852]
- 28 Iijima K, Otake T, Yoshinaga J, Ikegami M, Suzuki E, Naruse H, Yamana-oka T, Shibuya N, Yasumizu T, Kato N. Cadmium, lead, and selenium in cord blood and thyroid hormone status of newborns. *Biol Trace Elem Res* 2007;**119**:10-8. [17914214] [doi:10.1007/s12011-007-0057-1]
- 29 Ozdemir HS, Karadas F, Pappas AC, Cassey P, Oto G, Tuncer O. The selenium levels of mothers and their neonates using hair, breast milk, meconium, and maternal and umbilical cord blood in van basin. *Biol Trace Elem Res* 2008;**122**:206-15. [18301869] [doi:10.1007/s12011-008-8088-9]
- 30 Rhead WJ, Schrauzer GN, Saltzstein SL, Cary EE, Allaway WH. Vitamin E, selenium, and the sudden infant death syndrome. *J Pediatr* 1972;**81**:415-6. [5042504] [doi:10.1016/S0022-3476(72)80346-9]
- 31 Rayman MP. Food-chain selenium and human health: Emphasis on intake. *Br J Nutr* 2008;**100**:254-68. [18346308] [doi:10.1017/S0007114508939830]
- 32 Lorenzo-Alonso MJ, Bermejo-Barrera A, Cocho de Juan JA, Fraga Bermudez JM, Bermejo-Barrera P. Selenium levels in related biological samples: human placenta, maternal and umbilical cord blood, hair and nails. *J Trace Elem Med Biol* 2005;**19**:49-54. [16240672] [doi:10.1016/j.jtemb.2005.07.006]
- 33 Makhoul IR, Sammour RN, Diamond E, Shohat AI, Tamir A, Shamir R. Selenium concentrations in maternal and umbilical cord blood at 24-42 weeks of gestation: basis for optimization of selenium supplementation to premature infants. *Clin Nutr* 2004;**23**:373-81. [15158301] [doi:10.1016/j.clnu.2003.08.004]
- 34 Lee AM, Huel G, Godin J, Hellier G, Sahuquillo J, Moreau T, Blot P. Inter-individual variation of selenium in maternal plasma, cord plasma and placenta. *Sci Total Environ* 1995;**150**:119-27. [7878445] [doi:10.1016/0048-9697(95)04123-1]
- 35 Rudolph N, Wong SL. Selenium and glutathione peroxidase activity in maternal and cord plasma and red cells. *Pediatr Res* 1978;**12**:789-92. [693120] [doi:10.1203/00006450-197807000-00010]
- 36 Hostetler CE, Kincaid RL. Maternal selenium deficiency increases hydrogen peroxide and total lipid peroxides in porcine fetal liver. *Biol Trace Elem Res* 2004;**97**:43-56. [14742899] [doi:10.1385/BTER:97:1:43]
- 37 Dylewski ML, Mastro AM, Picciano MF. Maternal selenium nutrition and neonatal immune system development. *Biol Neonate* 2002;**82**:122-7. [12169835] [doi:10.1159/000063088]